

# AUDIT





## Introduction

The Alcohol Use Disorders Identification Test (AUDIT) is a 10-item screening tool developed by the World Health Organization (WHO) to assess alcohol consumption, drinking behaviors, and alcohol-related problems. Answer the AUDIT questions in terms of standard drinks. A chart illustrating the approximate number of standard drinks in different alcohol beverages is included for reference. A score of 8 or more is considered to indicate hazardous or harmful alcohol use. The AUDIT has been validated across genders and in a wide range of racial/ethnic groups and is well-suited for use in primary care settings. Detailed guidelines about use of the AUDIT have been published by the WHO and are available online: [http://whqlibdoc.who.int/hq/2001/who\\_msd\\_msb\\_01.6a.pdf](http://whqlibdoc.who.int/hq/2001/who_msd_msb_01.6a.pdf)

## The Alcohol Use Disorders Identification Test: Self-Report Version

PATIENT: Because alcohol use can affect your health and can interfere with certain medications and treatments, it is important that we ask some questions about your use of alcohol. Your answers will remain confidential so please be honest. Place an X in one box that best describes your answer to each question.

Questions	0	1	2	3	4	
1. How often do you have a drink containing alcohol?	Never	Monthly or less	2-4 times a month	2-3 times a week	4 or more times a week	
2. How many drinks containing alcohol do you have on a typical day when you are drinking?	1 or 2	3 or 4	5 or 6	7 to 9	10 or more	
3. How often do you have six or more drinks on one occasion?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
4. How often during the last year have you found that you were not able to stop drinking once you had started?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
5. How often during the last year have you failed to do what was normally expected of you because of drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
6. How often during the last year have you needed a first drink in the morning to get yourself going after a heavy drinking session?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
7. How often during the last year have you had a feeling of guilt or remorse after drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
8. How often during the last year have you been unable to remember what happened the night before because of your drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
9. Have you or someone else been injured because of your drinking?	No		Yes, but not in the last year		Yes, during the last year	
10. Has a relative, friend, doctor, or other health care worker been concerned about your drinking or suggested you cut down?	No		Yes, but not in the last year		Yes, during the last year	
					<b>Total</b>	

STANDARD DRINK EQUIVALENTS	APPROXIMATE NUMBER OF STANDARD DRINKS IN:
<b>BEER or COOLER</b>	
<p><b>12 oz.</b></p>  <p><b>~5% alcohol</b></p>	<p>12 oz. = 1  16 oz. = 1.3  22 oz. = 2  40 oz. = 3.3</p>
<b>MALT LIQUOR</b>	
<p><b>8-9 oz.</b></p>  <p><b>~7% alcohol</b></p>	<p>12 oz. = 1.5  16 oz. = 2  22 oz. = 2.5  40 oz. = 4.5</p>
<b>TABLE WINE</b>	
<p><b>5 oz.</b></p>  <p><b>~12% alcohol</b></p>	<p>a 750 mL (25 oz.) bottle = 5</p>
<b>80-proof SPIRITS (hard liquor)</b>	
<p><b>1.5 oz.</b></p>  <p><b>~40% alcohol</b></p>	<p>a mixed drink = 1 or more*  a pint (16 oz.) = 11  a fifth (25 oz.) = 17  1.75 L (59 oz.) = 39</p> <p>*Note: Depending on factors such as the type of spirits and the recipe, one mixed drink can contain from one to three or more standard drinks.</p>